AGRINTER-AGRIS

Total Control Control

TOTAL CONTROL

TOTAL

TOTA

PROGRAMME OF WORK
FOR THE SHORT-TERM ADAPTIVE
PRODUCTION ORIENTED RESEARCH
ON RICE IN THE
BRUMDEC PROJECT


IIGA PM-316 IICA/JAMAICA
Miscellaneous Publication #316
ISSN-0534-5391

FOR THE SHADOWARD RESEARCE OR SHEET THE LARD WARD WARD RESEARCE THE THE PROPERTY OF THE PROPER

Miscellaneous Phylocation =316



PROGRAMME OF WORK FOR THE SHORT-TERM ADAPTIVE PRODUCTION ORIENTED RESEARCH ON RICE IN THE BRUMDEC PROJECT



Vivian Chin
Rice Production Specialist
IICA/Jamaica

)

)

With $\mu_{i} = 0$, $\mu_{i} = 0$

00000433

The state of the s

1. INTRODUCTION

A review of the current state of technology in rice production and adaptive research on rice in the BRUMDEC Project area has been completed. (1)

On the basis of that review this short-term programme for adaptive production-oriented research on rice in the BRUMDBC Project is proposed.

This programme aims at correcting deficiencies by the transfer of technology and by generating information to enable the formulation of improved norms in the management of the rice crop.

2. OBJECTIVES

This programme of work has as its objectives:

- 2.1. Establishing the use of appropriate technologies that will correct observed imbalances in the rice production system on the BRUMDEC farm within the next two seasons.
- 2.2. Establishing fertilizer recommendations for rice grown on the Clay soils in the Project area.
- 2.3. Developing a complete technological package for rice production on the Morass Peat soil.
- 2.4. Byaluating the economics of rice production on each of the major rice soils in the Project area.
- 2.5. Training of National Technicians consistent with the overall agronomy of rice production.
- 2.6. Accumulating information that will provide the basis for planning of a longer-term programme of adaptive production-oriented research on rice and validation of the results and achievements attained.

in the composition of a precision of the community of the

in the company of the second control of the

2. The interpretation of the control of the cont

in the control of the

- The confidence of the confiden
- of the second second of the se
 - om province of general general sources of the source of th
- (A) a subject to the second of the second
- The complete distance of the complete c
 - The state of the s

3. STRATEGY

The strategy will be as follows:

3.1. In order that the programme produces immediate benefits, the identifiable deficiencies in the rice production system which can be corrected by applying appropriate methods will be given high priority for attention. The methods to be adopted have been dealt with in some detail in the review paper. (1)

Attention will be given to improving the quality of seed of the variety CICA 9, and any other variety needing such attention, up to internationally recognised quality standards.

- 3.2. Micro-and macro-plot experiments will be carried out to establish what should be the fertilizer practice for rice grown on the Clay soils and on the Morass Peat soil.
- 3.3. Micro and macro-plot experiments will be carried out on the main Clay soil -- Cashew Clay -- and on the Morass Peat soil to generate information for the formulation of a technology package for the successful cultivation of rice on those soils.
- 3.4. Variety x fertilizer trials will be carried out using varieties of rice developed in Colombia (CIAT-ICA), Dominican Republic, Guyana, Philippines (IRRI), Suriname and the United States of America. The varieties will be tested simultaneously at different levels of fertilization.

The objective of this part of the programme is to identify one or more varieties having a high level of adaptation to the environmental conditions of the Project area. All of the varieties to be tested will have rice quality (appearance and cooking characteristics) acceptable to the consumers.

3.5. In addition to the field studies planned for 3.2 and 3.3 above, pot trials will be carried out to assess the status of the micronutrient requirements of the rice plant when grown on soils of the Project area:

.

The first of the f

And the second of the second of

A Control of the Contro

- on Clay soils from fields which have not been cultivated with rice;
- on similar soils from fields which have been cultivated with rice for several seasons,
- on the Morass Peat soil which will be tested with rice for the first time.
- 3.6. Two training programmes will be conducted, one at the end of each season, for National technicians and farmers. The training programmes will embrace all important aspects of the agronomy of rice production.

4. METHODOLOGY FOR EXPERIMENTS, PRODUCTION STUDY, SEED PURIFICATION, AND TRAINING

4.1. Experimental Methods

4.1.1. Experimental design of the micro-plot studies

The design of the micro-plot experiments will be a Randomized Complete Block with two replications. Gross individual plot size will be 2.0 m x 1.0 m, with adjacent plots separated by an unplanted space of 0.5 m on all sides. Each micro-plot experiment will occupy 399 m². The effects of four (4) factors will be investigated:

- nitrogen at 0 kg/ha; 56 kg/ha, and 112 kg/ha.
- phosphate at 0 kg/ha, 28 kg/ha, and 56 kg/ha.
- potash at 0 kg/ha, 30 kg/ha; and 60 kg/ha.
- micro-nutrient mixture of chelated iron, chelated zinc, and chelated manganese, containing 3.12% of each of those elements, at 4.0053 kg mixture/ha, 8.0106 kg mixture/ha, and 12.0159 kg mixture/ha as foliar sprays at 35 days after sowing or earlier if necessary.

4.1.2. Experimental design of the macro-plot studies

A CONTROL OF THE CONT

The second of th

- Same and the second
- The state of the s

Compared to the second of the

La Contraction of the Contractio

4.1.2.1. Studies of NPK requirements of rice on soils of the BRUMDEC Project

The design of these studies on all soil types will be a $3 \times 3 \times 3$ NPK factorial with partial confounding of the NPK interaction, with two replicates having three blocks of nine plots each per replicate.

In these experiments the basal rate of application of micro-nutrients as foliar sprays will be:

- on the Clays: zinc at 0.25 kg/ha iron at 0.25 kg/ha, and manganese at 0.25 kg/ha.
- on the Morass Peat: zinc at 0.375 kg/ha; iron at 0.375 kg/ha, manganese at 0.375 kg/ha, and copper at 0.6 kg/ha.

The gross individual plot size will be 12 m x 7 m and the net plot size (harvested plot size) 10 m x 5 m (1/200th hectare). A border of 30 cm width contiguous to the net plot will be maintained intact. Also to be maintained intact is the outermost 30 cm wide border of the gross plot. The space (40 cm width) between the inner and outer border will be the sampling space for soil and plant samples to be taken during the course of the studies.

The treatment levels for the three factors will be:

- on the Clays: N at 0, 56 and 112 kg/ha. $P_2^0_5$ at 0, 28, and 56 kg/ha. K_2^0 at 0, 30, and 60 kg/ha.

The rate and timing of fertilizer will be:

tion

- nitrogen: 1/3 at 21 DAS*; 1/3 at panicle differentiation, and 1/3 at first heading.

- phosphate: ½ at rowing, and ½ at panicle differentiation

- potash: $\frac{1}{2}$ at 21 DAS; and $\frac{1}{2}$ at panicle differentia-

^{*} DAS = Days after sowing of germinated seed

The first the second of the second second

1. State of the second section of the second section is a second section of the second section of the second section is a second section of the section of the second section of the section of the second section of the section XX Commence of the commence of

the straining of the first of the second section is and the second of the second o

 $(-1)^{2} + (-1)^{2}$

The state of the s A Committee Comm

 $(-1)\nabla_{x}(\mathbf{G}_{x}^{-1}(x)) = (-1)^{2} \nabla_{x}(\mathbf{G}_{x}^{-1}(x)) + (-1)^{2} \nabla_{x}(\mathbf{G}$ The forest of the second section of the second $\mathbb{E}\left(\operatorname{res}_{\mathcal{A}}(\mathcal{A}^{\mathsf{p}}(\mathcal{A})) \otimes \mathbb{E}\left(\operatorname{res}_{\mathcal{A}^{\mathsf{p}}(\mathcal{A})} \otimes \operatorname{res}_{\mathcal{A}^{\mathsf{p}}(\mathcal{A$ energy of the control recording to the second of the

the second of th A STATE OF THE STA But the second of the second o

2 1, 1

A CARLO DE TRANSPORTE DE LA CARLO DEL CARLO DE LA CARLO DEL CARLO DE LA CARLO DEL CARLO DEL LA CARLO DEL LA CARLO DEL LA CARLO DE LA CARLO DEL LA CARLO DE LA CARLO DE LA CARLO DEL LA

reconstruction of

y a contract of the second : \, £

(x,y,y) = (x,y) + (y,y) + (y.

Cathering Company of the Company : .

- on the Morass Peat:

N at 0, 20 and 40 kg/ha. P_2^{0} at 0, 30, and 60 kg/ha. K_2^{0} at 0, 30 and 60 kg/ha.

The fertilizer application will be split in the same proportions as indicated for the treatments on the Clay soils.

These experiments will be repeated in the second season starting from October - November 1981.

4.1.2.2. Studies on the managment of tillage, irrigation water, and fertilizers in rice on the Cashew Clay soil and the Morass Peat soil

The design of this study on each soil type will be a Randomized Complete Block Split-Split-Plot Design with two (2) replicates.

Main plots:

Tillage management

Sub-plots:

Water management

Sub-sub-plots:

Fertilizer management

Factor definition:

- Tillage management 1 (T1): Dry tillage to produce a fine tilth, and levelling under dry conditions.
- Tillage management 2 (T2): Dry tillage initially to reduce
 the size of largest aggregates
 to approximately 7.5 cm diameter,
 flooding to an average depth of
 10 cm, puddling, under-waterlevelling;
- Tillage management 3 (T3): Flooding to an average depth of 10 cm, rototilling, and underwater-levelling.
- Water management. When tillage operations are completed under dry conditions all plots are to be submerged.

 All tillage on dry-tillage treated plots must

the second of th

•

the state of the s Reference of the second of the second

A Strain Committee of the Committee of t

Manager to the control of the contro and the second second

Control of the Contro

therefore be completed before flooding. Wet tillage operations will begin immediately after flooding and all plots will be kept submerged until two (2) days after all wet-tillage operations are completed.

At two (2) days after the completion of wet-tillage operations the following water management treatments will begin:

- Water management treatment 1 (W1):

- Retain average depth of 7.5 cm flood;
- Seal all levees; allow seedlings to emerge through flood;
- Irrigate to renew 7.5 cm average depth of flood when average depth of flood reduces to 2.5 cm,
- When plants are approximately 14 cm tall, increase depth of flood to 10 cm and renew to this depth when flood depth reduces to 2.5 cm.
- Drain plot completely at 15 days after 50% heading.
- Water management treatment 2 (W2)
 - Retain average depth of 7.5 cm flood (by draining off excess water if necessary).
 - Sow seed
 - Seel all levees, do not irrigate (if rainfall increases flood depth, drain off water to maintain 7.5 cm depth of flood)
 - Grow seedlings through this depth of flood and maintain daily records of depth of flood in each plot.
 - Irrigate to renew 7.5 cm average depth of flood when the soil surface becomes exposed to the atmosphere.

and the second of the second o

Tark

- Allow evapotranspiration, percolation and seepage to reduce flooding depth until the soil surface is just exposed, then irrigate to 10 cm average depth of flood.
- Continue this sequence of topping up plot to 10 cm average depth of flood when the soil surface of plot is just exposed.
- Drain plot completely at 15 days after 50% heading.

- Water management treatment 3 (W 3)

- Drain plot completely.
- Sow Seed
- Keep open all plot-drainage outlets from plot to allow rainfall water to exit.
- All 10 DAS irrigate to a depth of 75% seedling height. Record the average depth of flood in each plot.
- Irrigate to increase average depth of flood to 10 cm. When the average depth of flood reduces to 2.5 cm.
- Continue this sequence of topping up plot to 10 cm average depth of flood when the average depth reduces to 2.5 cm.
- Drain plot completely at 15 days after 50% heading.

- Water management treatment 4 (W 4)

- Drain plot completely
- Sow Seed
- Keep open all plot drainage outlets to allow rainfall water to exit.
- At 10 DAS irrigate to a depth of 75% seedlings height. Record the average depth of flood in each plot.

The state of the s

$\mathcal{L}_{i} = \mathcal{L}_{i} + \mathcal{L}_{i}$

. :

 $(1+\epsilon)^{2} + (1+\epsilon)^{2} + (1+\epsilon$

and the second of the second of the second

Superior of the s

- Allow evapotranspiration percolations and seepage to reduce flood depth until the soil surface of the plot is just exposed then irrigate to 10 cm average depth when the soil surface of the plot is just exposed.
- Drain plot completely at 15 days after 50% heading.

4.1.3. Fertilizer Management Treatments

There will be seven (7) fertilizer management treatments:

4.1.3.1. On Cashew Clay Soil

Fertilizer Management Treatment	Kg/Ha		Rate of Timing of Fertilizer
FMCCO	$\frac{N}{2} \frac{P_2^0_2}{P_2^0_2}$		None
FMCC1	0 0 64.4 28.8 3	0 6.6	28.4 Kg N/ha + 28.8 P ₂ 0 ₅ /ha + 33.6 Kg K ₂ 0/ha just prior to sowing
FMCC2	64.4 28.8 3	3.6	12 Kg N/ha at 21 DAS 24 Kg N/ha at panicle differentiation 12 Kg N/ha + 28.8 Kg
			P ₂ 0 ₅ /ha + 33.6 Kg K ₂ 0/ha just prior to sowing 28.4 Kg N/Ha at 21 DAS 24 Kg N/ha at panicle differentiation
FMCC3	64.4 28.8 3	3.6	12 Kg N/ha + 14.4 Kg P ₂ 0 ₅ /ha + 16.8 Kg K ₂ 0/ha just prior to sowing 12 Kg N/ha at 21 DAS 24 Kg N/ha + 14.4. Kg
			P ₂ 0 ₅ + 16.8 Kg K ₂ 0/ha at panicle differentia- tion

. . . .

and the second of the second		1 •
· :		
••		
1 · · .	· ·	•
• .•		
		•
	<u>.</u>	
, · · · · ·		
6.2		
,		

• • • •

Fertilizer Management Treatment	Kg/H	la	Rate and Timing of Fertilizer
FMCC4	N P ₂ 0 ₅ 64.4 28.8	K ₂ 0 33.6	12 Kg N/ha + 14.4 kg P ₂ 0 ₅ /ha + 16.8 kg K ₂ 0/ha just prior 28.4 Kg N/ha at 21 DAS 24 Kg N/ha + 14.4 Kg P ₂ 0 ₅ /ha at panicle differentiation
FMCC5	50 25	25	10 Kg N/ha + 12.5 Kg P ₂ 0 ₅ /ha + 12.5 Kg K ₂ 0/ha just before sowing. 20 Kg N/ha at 21 DAS 20 Kg N/ha + 12.5 Kg P ₂ 0 ₅ + 12.5 Kg K ₂ 0/ha at panicle differentiation.
FMCC6	50 25	25	10 Kg N/ha + 25 Kg P ₂ 0 ₅ /ha + 25 Kg K ₂ 0/ha just before sowing. 20 Kg N/ha at 21 DAS 20 Kg N/ha at panicle differentiation.

.

e e e e e e e e e e e e e e e e e e e			· · · · · · · · · · · · · · · · · · ·
	•		•
		•	
•			
t ,			
. t	•		
et en			
	•		
\$2.	,	•	
$\mathbf{e}_{i} = \mathbf{e}_{i} \mathbf{e}_{i}$			
,	•		
	•	1	
		:	
party Notice (1911)		•	
State of the state	i	,	• 1
		!	
A Committee of the Comm			
		•	
· · · · · · · ·			• • •

On Morass Peat Soil	ŭ	Kg/H	la .	Rate and Timing of Fertilizations
FMMP 0 FMMP 1	0 10	0 10	0 20	None 10 Kg P ₂ 0 ₅ /ha) 20 Kg K ₂ 0/ha) 10 Kg N/ha)
FMMP 2	10	10	20	10 Kg P ₂ 0 ₅ /ha) 20 Kg K ₂ 0/ha)at sowing 10 Kg N/ha) at P.D.
FMMP 3	20	20	30	10 Kg P ₂ 0 ₅ /ha) 20 Kg K ₂ 0/ha)at sowing 10 Kg P ₂ 0 ₅ /ha) 10 Kg K ₂ 0/ha)at P.D.
FMMP 4	20	20	30	20 Kg N/ha
FMMP 5	30	30	40	20 Kg K ₂ 0/ha) at sowing 20 Kg P ₂ 0 ₅ /ha) 20 Kg K ₂ 0/ha) 10 Kg P ₂ 0 ₅ /ha) at P.D.
FMMP 6	30	30	30	30 Kg N/ha) 40 Kg K ₂ 0/ha) 20 Kg P ₂ 0 ₅ /ha) at sowing 10 Kg P ₂ 0 ₅ /ha) 30 Kg N/ha) at P.D.

In these rice crop management experiments basal micro-nutrient fertilization will be at the rates indicated for the Clays and Morass Peat in 4.1.2.1. above.

. •

These experiments will be repeated in the second season beginning October - November 1981.

4.1.2.3 Variety Trials

In this season's work only four varieties will be included in the variety trials. The varieties will be CICA 8, CICA 9, Variety "N" and Rustic. The first three named varieties are long-grained. Rustic has an extra long grain.

On the matter of consumer acceptance the Jamaican public has been supplied with rice of variety "N" and Rustic in rice shipments imported from Guyana.

In the second season's work it is planned to include in addition to the four varieties named above, one variety from the Dominican Republic (JUMA 57), two varieties from Suriname (ELONI and Diwani) another variety from Guyana (Champion), and three varieties from the U.S.A. (Bluebelle, Starbonnett, Lebonnet).

The design of the first season's variety trial has been changed to permit the testing of the four varieties at different levels of fertilization. The design will be a Latin Square Split Plot with varieties occupying the main plots and fertilizer levels occupying sub-plots. This study will be done on Cashew Clay and Morass Peat Soil.

The s	sub-plot	treatments	will	be as	follows:
-------	----------	------------	------	-------	----------

Fertilizer Level	N	Kg/ P	Ha K	Rate and Timing of Fertilization
CCF 0	0	0	0	None
CCF 1	56	20	25	10 Kg P ₂ 0 ₅ /ha) 15 Kg K ₂ 0/ha) at sowing
				28 Kg N/ha at 21 DAS at panicle 28 Kg N/ha differentiation 10 Kg P ₂ 0 ₅ /ha) 10 Kg K ₂ 0/ha 29 Kg P ₂ 0 ₅ /ha) 34 Kg K ₂ 0/ha 28 Kg N/ha at 21 DAS
CCF 2	64	29	34	29 Kg P ₂ 0 ₅ /ha) at sowing 34 Kg K ₂ 0/ha) 28 Kg N/ha at 21 DAS 12 Kg N/ha at 42 DAS 24 Kg N/ha at panicle differentiation

en de la companya de la co

The second of th

 $(1.86 \pm 0.00) \times (1.16 \pm 0.00) \times (1.00) \times (1.00$

2017 1114					or the state of the state of
			, practions	•	
	<i>;</i>	<i>;</i> •			7.
	$0 \leq \frac{1}{2} \leq \frac{2}{3} \leq \frac{1}{3}$	i		;	
;	· · · · · · · · · · · · · · · · · · ·			i	
		. :		1	
		•		,	,
				!	

On Morass	Peat Soil	N	Kg/ P	Ha K	Rate and Timing of Fertili- zation
MPF	0	0	0	0	None
MPF	1	10	10	20	10 Kg P ₂ 0 ₅ /ha)at sowing 20 Kg K ₂ 0/ha) 10 Kg N/ha at panicle differentiation
MPF	2	20	20	30	10 Kg P ₂ 0 ₅ /ha) 20 Kg K ₂ 0/ha)at sowing 10 Kg P ₂ 0 ₅ /ha)at panicle 10 Kg K ₂ 0/ha)differentia- 20 Kg N/ha)

For the variety X fertilizer trials the micro-nutrient fertilization levels on the Clay soil and on the Morass Peat soil will be as indicated in 4.1.2.1. above.

4.1.2.4. Pot Trials

The pot trials for the first season's work will test the response of the rice plant to foliar applications of selected micronutrients. The ph ranges of the Clays and the Morass Peat (4) suggest that minor element nutrition should be investigated.

The minor elements to be used are:-

On Clay Soils:

- Iron, manganese and zinc

On the Morass Peat Soil:

- Iron, manganese, zinc and copper

The design of the experiments will be:

- on the Cashew Clay and Wallen's Clay 2⁸ factorial in a completely Randomized Design with two replicates.

A control of the contro

The state of the s

The state of the second second

 $\label{eq:condition} (c) = (c + 1)^{n} + ($

The state of the s

 $\frac{1}{2}\left(\frac{1}{2}\right)\right)\right)}{\frac{1}{2}}\right)\right)}\right)\right)}\right)}\right)}\right)}\right)}\right)}\right)}\right)}$

 on the Morass Peat Soil: 2⁴ factorial in a completely Randomized Design with two replicates.

Basal NPK fertilization will be applied as follows:

- on the Clay soils:

60 Kg N/ha

30 Kg P₂0₅/ha

30 Kg K_2 0/ha

- on the Morass Peat Soil

30 Kg N/ha

 $30 \text{ Kg P}_{2}0_{5}/\text{ha}$

40 Kg K₂0/ha

4.1.2.5. Rice Production Study

On the Cashew Clay Soil type a state of the art production plot of ten (10) hectares (24.71 acres) will be managed as per a prorated budget for 10 hectares. A budget for 405 hectares has been presented earlier. (1) Actual costs and returns will be recorded.

4.1.2.6. Seed Purification

Seed purification plots of all varieties entered in the variety/fertilizer trial will be grown in the first season. The entire area for seed purification work will not exceed one (1) hectare (2.471 acres). Separate nurseries will be prepared for each variety. Each variety will occupy approximately 0.25 hectare (0.618 acres) in the field after transplanting. Spacing will be 30 cm x 15 cm at one (1) seedling per hill.

We expect to obtain approximately 800 kg. (1,763 lbs) of bulked seed and 100 single-plant selections of each variety after purification in the field is completed.

4.1.2.7. Training Programme

On the Cashew clay and Morass Peat soil types, demonstration plots of size 0.4 hectare (approximately 1 acre) will be sown with variety CICA 9 at 14 day intervals. There will be ten (10) sowing dates on each soil type.

The training programme of each season will be planned to begin on DAY 119 and to end on DAY 126.

,

. . .

and the Substitute of the Control of

Here is the state of the state

A control of the contro

$\sum_{i=1}^{n} \frac{\partial u_i}{\partial x_i} = \sum_{i=1}^{n} \frac{\partial u_i}{\partial x_i} = \sum_{i=1}^{n} \frac{\partial u_i}{\partial x_i} = \sum_{i=1}^{n} \frac{\partial u_i}{\partial x_i} = 0$

A fine of the control o

Participants will be exposed to the rice crop at all important stages of its development from land preparation through to harvesting.

Field work will occupy one-half of each training day and class room lectures will be conducted for not more than 1 hour each day. Some time will be allocated each day for participants to familiarise themselves with the different types of agricultural chemicals used in rice production.

4.1.3. Data Collection

4.1.3.1. Soil Analysis

To monitor nutrient status of the soil prior to each fertilizer application and at maturity.

Sampling at: prior to 1st tillage; 21 DAS, 63 DAS, 126 DAS

Sampling Date	Type:Sample	To be analysed for
Prior to 1st tillage	Soil Sample	% Clay, % silt, % sand % organic carbon P ₂ 0 ₅ - ppm (Truog) K - m.e./100g soil Na - m.e./100g soil Mg - m.e./100g soil Ca - m.e./100g soil SO ^S - m.e./100g soil, non
		Sulphare - S C1 ⁴ - me./100g soil Mn - ppm Zn - ppm Fe - ppm B - ppm Mo - ppm Cu - ppm
AT 21 DAS, 63 DAS	Soil Sample	% organic carbon P ₂ 0 ₅ - (ppm) (Truog) K - m.e./100g soil Na -m.e./100g soil; Mg - m.e./100g soil SO ⁵ -m.e./100g soil; non Sulphare - S C1 ⁴ - m.e./100g soil Mn - ppm, Fe - ppm, B- ppm Zn - ppm; Mo - ppm

and the

 $F_{ij} = F_{ij} = F_{ij}$

•

•

1

٠.

.

4.1.3.2. Plant tissue analysis

At 21 DAS, 42 DAS 63, DAS, Plant Dry weight/100 cm² plot area, 84 DAS, 105 DAS, 126 DAS. Sample content of N,P,K, Ca, Na, Mg, SO₄, Cl, Mn, Zn, Fe, and B in plant tissue.

- 4.1.3.3. Grain (at 12% moisture) (g) / Straw (Dry weight) (g) ratio at maturity (126 DAS).
 - From two (2) 10 cm x 10 cm. Sub-samples in micro plot studies
 - From four (4) 30 cm x 30 cm.
 Sub-samples in macro-plot studies.
- 4.1.3.4. Grain yield will be recorded only from macro-plot studies on the basis of kg/ha of clean grain at 12.5% moisture.
- 4.1.3.5. Data on number of tillers per unit area will be recorded at the same time when plants and soil are sampled for chemical analyses.
- 4.1.3.6. Plant height of 10 randomly selected plants/plot at maturity will be recorded to the nearest centimetre. Plant height will be measured from the soil surface to the tip of the panicle of the tallest tiller each randomly selected plant (excluding awns if present).
- 4.1.3.7. The following ancillary data will be recorded from the samples taken at 4.1.3.3. above.
 - Number of grains/panicle
 - Weight of 100 grains (g)
- 4.1.3.8. From the grain harvested to provide yield data in the microplot studies, two 500g samples of clean dried grain at a moisture content of 12.5% for each plot will be dehulled using a laboratory shelter and milled in a laboratory mill for 30 seconds. The data to be collected will be:
 - weight of dehulled sample (g) and by difference find weight of hulls (g) and calculate % by weight of hulls.
 - weight of milled rice, and calculate % total recovery.

en de la companya de la co

en de la companya de la co

- weight of head rice (g) and weight of brokens (g) separately, and calculate % recovery of head rice and % recovery of brokens.
- by difference between weight of dehulled sample (g) and weight of milled rice (g) find weight of bran and calculate % recovery of bran.
- Percentage by weight of chalky grains in the hulled rice sample. A chalky grain for the purpose of this assessment is defined as a rice grain exhibiting one or more of the following blemishes:
 - (a) white belly
 - (b) white centre
 - (c) white back

A random sub-sample of the chalky grains will be assessed for the extent of chalkiness using an internationally accepted scale. (3)

Table I: Scale for Opacity of the Rice Endosperm (3)

Scale	Chalkiness (expressed as % Surface Area)
0	None (0%)
1	Low (less than 10%)
5	Medium (10% to 20%)
9	High (more than 20%)

4.1.3.9. Condition of Crop

Specific notes will be made of the condition of the crop in the field from sowing through harvesting with reference to lodging of the crop, the occurrence of pests, weeds, and diseases, the type of control measures used and the effect of such control measures.

•

٠,

.

4.1.3.10. Meteorological Data

It is desirable that during the cause of the experiments we should be able to obtain daily records of the following meteorological data:

- rainfall (mm)
- maximum and minimum temperatures (°C)
- relative humidity at 0900 hours and at 1400 hours
- Solar radiation () or if such data are not available, number of hours of sunshine per day
- open pan evaporation rate.

4.1.3.11 Quality of Irrigation Water

Routine sampling of irrigation water entering experimental blocks will be done at each irrigation time-run. Samples will be sent to a laboratory to be analysed for:

- electrical conductivity
- bicarbonate content
- pH

4.1.3.12. Soil Moisture Prior to Dry Tillage Operations

Routine tests will be done to obtain data.

ADDITIONAL STUDIES IN SECOND SEASON OF WORK

All main studies done in the first season will be repeated in the second season to test the repeatability of the results. From the results of the first season's work the best treatment or best treatment-combination will be used as the median level and will be bracketed by lower and higher treatment levels.

All factors of which a treatment level has been adjudged to be significantly better than the other levels tested will be included in a separate study to be done on the Cashew Clay soil and the Morass Peat. This additional study will provide in greater detail, information on the response surface.

± ...

. . . .

the state of the s

tana and tanàna mpikambana ao amin'ny fivondronana ao amin'ny fivondronana ao amin'ny fivondronana ao amin'ny Ny INSEE dia mandronana ao amin'ny faritr'i Austria ao amin'ny faritr'i Austria ao amin'ny faritr'i Austria ao

in the time is a first of the contract of the

e e e

 $((a_1,\ldots,a_{n-1}),(a_1,\ldots,a_{n-1}),(a_1,\ldots,a_{n-1}))$

the state of the s

REFERENCES

1. CHIN, A.V.E. 1981,

Rice research and production in the BRUMDEC Project. State-of-the-art- review, identification of constraints and interim recommendation and budget for establishing 4 or 5 hectares (1,000 acres) of rice on the clay soils at BRUMDEC. IICA, Kingston, Jamaica - June 1981 29p.

2. WAHAB ABDUL H. (undated)

Determination of the nutritional status of soils by means of the micro-plot technique.

Inter-American Institute for Co-operation on Agriculture - OAS, Jamaica Office, Kingston, Jamaica 9p.

3. CENTRO INTERNACIONAL DE AGRICULTURA TROPICAL (CIAT) (undated)

Sistema de Evaluacion Estandar para Arroz CIAT, Cali., Colombia 62p.

4. HEWIT C.W. 1978.

Report on the detailed Soil Survey of the Upper Morass. Black River Upper Morass Development Company Limited (BRUMDEC). March 1978. 16p Tables.

200

The control of the state of the

11

I MAN TEST OF THE REPORT OF A SECTION OF THE SECTIO

The decay me material to be as a constant Colored Control of the Colored Colored Burney Barrier Barrier

The state of the s The state of the s and the first of the second

and the second of the second o Section 1995 to the section of the s and the second of the second o

Contract of the Contract of th

All the second of the second o The social state of the so College Control of the State of f₂ 1

AGRICULTURE IN JAMAICA

Collection of papers of the Office of IICA in Jamaica

1977 -	19	<u> 78</u>	
No. I	-	1	Fritz Andrew Sibbles, "Basic Agricultural Information on Jamaica Internal Document of Work", January 1977
No. I	-	2	Yvonne Lake, "Agricultural Planning in Jamaica", June 1977
No. I	-	3	Aston S. Wood, Ph.D., "Agricultural Education in Jamaica", September - October 1977
No. I	-	4	Uli Locher, 'The Marketing of Agricultural Produce in Jamaica', November 1977
No. I	-	5	G. Barker, A. Wahab, L. A. Bell, "Agricultural Research in Jamaica", November 1977
No. I	-	6	Irving Johnson, Marie Strachan, Joseph Johnson, "Land Settlement in Jamaica", December 1977
No. I	-	7	Government of Jamaica, "Agricultural Government Policy Papers", February 1978
No. I	-	8	Jose Emilio Araujo, "The Communal Enterprise", February 1980
No. I	-	9	IICA and MOAJ, "Hillside Farming Technology - Intensive Short Course", Vols. I and II, March 1978
No. I	- :	10	Jose Emilio Araujo, "The Theory Behind the Community Enterprise - Seminar in Jamaica", March 1978
No. I	- 1	1	Marie Strachan, "A National Programme for the Development of Hillside Farming in Jamaica", April 1978
No. I	- 13	2	D. D. Henry, "Brief Overall Diagnosis of Hillside Farming in Jamaica", April 1978
No. I	- 13	3	Neville Farquharson, "Production and Marketing of Yams in Allsides and Christiana", May 1978

:			(* * * * * * * * * * * * * * * * * * *	
	•			
the second of th			$\frac{s}{1} \leq \frac{s}{2} \leq \frac{s}{2}$	ĩ
				e se
			,	
	e en E	. :		;
	No. 10 Percentage of the control of	•	·	· .
				1
		,	winds in	,,,
		.:		** - 1

- No. I 14

 R. C. E. McDonald, A. H. Wahah, "Fertility Assessment of Newly Terraced Hillside Soils Using the Microplot Technique the Allsides Case Study", 1978
- No. I 15 IICA IDB, "Course in Preparation and Evaluation of Agricultural Projects", Vols. I and II, November 1977
- No. I 16 Neville Farquaharson, "Production and Marketing of Dasheen in Allsides and Christiana", June 1978

1978 - 1979

- No. II 1 O. Arboleda-Sepulveda (IICA-CIDIA), "Agricultural Documentation and Information Network in Jamaica", September 1978
- No. II 2 Victor Quiroga, "National Agricultural Information System", (NAIS-Jamaica) Project Profile, September 1978
- No. II 3 Joseph Johnson, "A Review on Land Reform in Jamaica for the Period 1972 1978", September 1978
- No. II 4 Neville Farquharson, "ABC of Vegetable Farming", A
 Draft High School Textbook, Vols. I, II, III and IV,
 February 1979
- No. II 5 Jerry La Gra, "Elements of an Agricultural Marketing Strategy for Jamaica", March 1979
- No. II 6 D. D. Henry, I. E. Johnson, "Agricultural Extension Service in Jamaica", March 1979

1979 - 1980

- No. III 1 H. R. Stennett, "Watersheds of Jamaica and Considerations for an Ordinal Scale of their Development", July 1979
- No. III 2 IICA-MAJ, 'Hillside Farming in Jamaica', A Training Seminar, December 1978
- No. III 3 A. L. Wright, A. H. Wahab, H. Murray, "Performance of Six Varieties of Red Peas (Phaseolus vulgaris L.)
 on a Newly Terraced Ultisol in Jamaica", September 1979
- No. III 4 IICA/Jamaica Staff, "Agro-Socio-Economic Sample Survey of Allsides Trelawny, Jamaica", September 1979

e de la companya de l

No. 111 - 5	Hilly Lands", October 1979
No. III - 6	IICA-MOAJ, "Tree Crops of Economic Importance to Hillside Farms in Jamaica", October 1979
No. III - 7	Canute McLean, "Production and Marketing of Peanuts", November 1979
1980	
No. IV - 1	Joseph Johnson, "Production and Marketing of Red Peas in the Hilly Areas of Jamaica", January 1980
No. IV - 2	Lyn Snuffer, "Rural Women: An Annotated Caribbean Bibliography with special reference to Jamaica", January 1980
No. IV - 3	Vincent Campbell, Abdul Wahab, Howard Murray, "Response of Peanut (Arachis hypogaea L.) on a Newly Terraced Ultisol in Jamaica", January 1980
No. IV - 4	P. Aitken, A. Wahab, I. Johnson, A. Sahni, "Agro-Socio-Economic Survey - Pilot Hillside Agricultural Project 'PHILAGRIP' Southern Trelawny", February 1980
No. IV - 5	Glenys H. Barker, "Bibliography of Literature relating to Research and Development in the Agricultural Sector of Jamaica 1959 - 1979", March 1980
No. IV - 6	Milton R. Wedderburn, "Allsides Farmers' Pre-Cooperative A Socio-Economic Assessment", March 1980
No. IV - 7	Adele J. Wint, "The Role of Women in the Development Process", April 1980
No. IV - 8	Milton R. Wedderburn, "The Co-operative Input in the Development of the Pilot Hillside Agricultural Project (PHILAGRIP)", April 1980
No. IV - 9	MOJ/IICA/CARDI, Fruit Trees Seminar - "Research & Development of Fruit Trees", June 1980
No. IV - 10	Henry Lancelot, 'Traditional Systems in Hillside Farming, Upper Trelawny, Jamaica', June 1980

ELICINE LEGISLA ELICINE PROPERTO DE LA CONTRA DEL CONTRA

- Vi IV
- - - (2) A Company of the Company of the Company of the Months of the Company of th

- No. IV 11 IICA/Jamaica, "Pilot Hillside Agricultural Project", (PHILAGRIP), Project Document. Vols. I, II and III, June 1980
- No. IV 12

 A. Wahab, I. Johnson, P. Aitken, H. Murray and
 H. Stennett, "Highlights of the Pilot Hillside
 Agricultural Project at Allsides", July 1980
- No. IV 13

 I. Johnson, A. Wahab, P. Aitken, H. Payne, "Benchmark for a Project Profile for Developing a Peanut Industry in Jamaica", July 1980
- No. IV 14 P. Aitken, A. Wahab, I. Johnson, "The Allsides Post Peasant", August 1980
- No. IV 15 Norma Munguia, Percy Aitken, Abdul Wahab, Irving Johnson, "Salt Extraction by Solar Energy", A Miniproject, September 1980
- No. IV 16

 Abdul H. Wahab, Percy Aitken-Soux, Irving E. Johnson and Howard Murray, "The Allsides Project in Jamaica Developmental Potentials of Hillside Agriculture", September 1980
- P. Aitken, A. Wahab, I. Johnson, A. Sahney and N. Munguia, "Rural Women Survey", Vols. I, II and III, October 1980
- No. IV 18
 P. Aitken, I. E. Johnson, A. Wahab, "Assessment of Employment Among Small Hillside Farmers of Jamaica", November 1980
- No. IV 19 IICA/Jamaica "Pilot Hillside Agricultural Project", (PHILAGRIP), Final Project Document. October 1980.
- No. IV 20
 P. Aitken, A. Wahab, I. E. Johnson, Bo-Myeong Woo,
 "IICA Evaluation of the First Phase FSB Allsides
 Project", (Internal Document of Work), November 1980
- No. IV 21 MINAG/IICA/CARDI "Seminar on Multiple Cropping", December 1980

No. V - 1

N. Munguia, P. Aitken, A. Wahab, I. Johnson, "Smoke

Curing of Fish (as a household industry in Rural Jamaica)",

January 1981

٠,
No. of the second
ىم
1
e. V
1
* * v

No. V - 2	P. Aitken, A. Wahab, I. Johnson, "Under-employment - It's Relation to the Agricultural Sector and Considera- tions for its Management", January 1981
No. V - 3	D. D. Henry, J. R. Gayle, "The Culture of Grafted Pimento (as spice crop for Allsides, Jamaica)", January 1981
No. V - 4	Abdul H. Wahab, Noel Singh, "Agricultural Research in Jamaica", February 1981
No. V - 5	P. Aitken-Soux, A. H. Wahab, I. E. Johnson, "Country Level Action Plan (CLAP)", May 1981
No. V - 6	P. Aitken-Soux, A. H. Wahab, I. E. Johnson, 'Overview of Agricultural Development in Jamaica', May 1981
No. V - 7	Samuel Thompson, I. E. Johnson, P. Aitken-Soux, Abdul Wahab, "The Land Development & Utilization Act 1966", July 1981
No. V - 8	Abdul Wahab, Percy Aitken-Soux, Irving Johnson, Bo-Myeong Woo, Howard Murray, Joseph Dehaney, "The Experiences of Jamaica in the Management of Agricultural Production on Hillsides", July 1981
No. V - 9	Dave Hutton, Abdul Wahab, Howard Murray, "Yield Response of Yellow Yam (Dioscorea Cayenensis) After Disinfesting Planting Material of Pratylenchus Coffeae", July 1981
No. V - 10	Elaine Montague-Gordon, Abdul H. Wahab, Joseph Dehaney and Audrey Wright, "Performance of Eleven Varieties of Dry Beans (Phaseolus vulgaris) Over Two Successive Seasons on the Hillsides of Jamaica", August 1981
No. V - 11	Dave G. Hutton, Abdul H. Wahab, "Position Paper on Root Crops in Jamaica", August 1981
No. V - 12	Percy Aitken-Soux, Abdul H. Wahab, Irving E. Johnson, "Technical Assistance for the English Speaking Caribbean (Considerations for an IICA Strategy)" (Internal Document of Work), September 1981
No. V - 13	Bo-Myeong Woo, Abdul H. Wahab, Joseph Dehaney, "Crop Production on Hillsides using non-Bench Terracing Alternative Measures for Soil Conservation (first year's results of the Olive River Soil Conservation studies)", September 1981

			•
	•	- \;	. w [.] .
	-	- •	
Application of the second of t	,		í
		٠	1
	-	•	; ;
	•		• •
	, •		
	, .		•
	<u>'</u>	- ··	•

No.	V - 14	Bo-Myeong Woo, Howard Murray and Joseph Dehaney, "Agricultural
		Production on Hillsides - the Allsides Project Case Study", September 1981
No.	V - 15	D. G. Hutton, A. H. Wahab and J. Dehaney, "Investigating Critical Levels of Dry Rotting of Yellow Yam (Dioscorea Cayenensis) Planting Material, the Benefits of Disinfesting the Heads of Pratylenchus Coffeae and of After-Planting Nematicide Treatments", September 1981
No.	V - 16	D. G. Hutton, A. H. Wahab, H. Murray and J. Dehaney, "Critical Levels of Dry Rotting of Yellow Yam (Dioscorea Cayenensis) Planting Material and Yield Responses After Disinfesting Heads of Pratylenchus Coffeae and After Post-Plant Nematicide Applications", September 1981
No.	V - 17	E. Ayer and J. Reyes, "Seminar on Mediterranean Fruit Fly", September 30, 1981
No.	V - 18	Bo-Myeong Woo, "Erosion Control Works in Korea", October 1981
No.	V - 19	Irving E. Johnson and Percy Aitken-Soux, "Country Level Action Plan (CLAP)" (Third Revision - Internal Document of Work), October 1981
No.	V - 20	Humberto Pizarro, "Programme of Work to Establish Guidelines for the Effective Administration, Operation and Maintenance of the Irrigation and Drainage District in the BRUMDEC Project November 1981
No.	V - 21	Humberto Pizarro, "The Operation of the Drainage System in the Black River Upper Morass Project", November 1981
No.	V - 22	Humberto Pizarro, "Recommendations for Land Use and Irrigation Needs in the BRUMDEC Project", November 1981
No.	V - 23	Humberto Pizarro, "Organization, Operations and Maintenance of the Irrigation System in the BRUMDEC Project", November 1981
No.	V - 24	Humberto Pizarro, "Basic Information for Planning Water Management in the BRUMDEC Project", November 1981

V

- No. VI 1

 Vivian Chin, "Rice Research and Production in the BRUMDEC

 Project State-of-the-Art Review, Identification of

 Constraints and Interim Recommendations and Budget for

 Establishing 405 Hectares (1,000 Acres) of Rice on the

 Clay Soils at BRUMDEC", January 1982.
- No. VI 2 Vivian Chin, "Programme of Work for the Short-Term Adaptive Production Oriented Research on Rice in the BRUMDEC Project", January 1982.

	•		EVOLUCIO	1	
,**					
·					
		ļ			
					:
		<u>J </u>	l	<u> </u>	1

•

.



DOCUMENTO
MICROFILMADO
Fecha: 15 NOV 1982